

Simulation of jet quenching at RHIC and LHC

I P Lokhtin, A M Snigirev

Skobeltsyn Institute of Nuclear Physics, Moscow State University, Moscow, Russia

E-mail: Igor.Lokhtin@cern.ch

Abstract. The model to simulate jet quenching effect in ultrarelativistic heavy ion collisions is presented. The model is the fast Monte-Carlo tool implemented to modify a standard PYTHIA jet event. The model has been generalized to the case of the “full” heavy ion event (the superposition of soft, hydro-type state and hard multi-jets) using a simple and fast simulation procedure for soft particle production. The model is capable of reproducing main features of the jet quenching pattern at RHIC and is applied to analyze novel jet quenching features at LHC.

PACS numbers: 25.75.-q, 12.38.Mh, 24.85.+p

1. Introduction:

One of the important tools for studying the properties of quark-gluon plasma (QGP) in ultrarelativistic heavy ion collisions is the analysis of a QCD jet production. The medium-induced energy loss of energetic partons, “jet quenching”, should be very different in the cold nuclear matter and QGP, resulting in many observable phenomena [1]. Recent RHIC data on high- p_T particle production at $\sqrt{s} = 200$ A GeV are in agreement with the jet quenching hypothesis [2]. At LHC, a new regime of heavy ion physics will be reached at $\sqrt{s_{NN}} = 5.5$ A TeV where hard and semi-hard particle production can stand out against the underlying soft events. The initial gluon densities in Pb+Pb reactions at LHC are expected to be much higher than those at RHIC, implying a stronger partonic energy loss, observable in new channels.

In the most of available Monte-Carlo heavy ion event generators the medium-induced partonic rescattering and energy loss are either ignored or implemented insufficiently. Thus, in order to analyze RHIC data on high- p_T hadron production and to test the sensitivity of LHC observables to the QGP formation, the development of adequate and fast Monte-Carlo tool to simulate the jet quenching is necessary.

2. Physics model and simulation procedure

The detailed description of physics model can be found in our recent paper [3]. The approach bases on an accumulating energy loss, the gluon radiation being associated with each parton scattering in the longitudinally expanding quark-gluon fluid and

includes the interference effect using the modified radiation spectrum dE/dl as a function of decreasing temperature T . The basic kinetic integral equation for the energy loss ΔE as a function of initial energy E and path length L has the form

$$\Delta E(L, E) = \int_0^L dl \frac{dP(l)}{dl} \lambda(l) \frac{dE(l, E)}{dl}, \quad \frac{dP(l)}{dl} = \frac{1}{\lambda(l)} \exp(-l/\lambda(l)), \quad (1)$$

where l is the current transverse coordinate of a parton, dP/dl is the scattering probability density, dE/dl is the energy loss per unit length, λ is in-medium mean free path. The collisional loss in high-momentum transfer limit, radiative loss in BDMS approximation [4], simple Gaussian parameterization of gluon emission angle distribution and realistic nuclear geometry are used. The QGP proper formation time τ_0 and initial temperature T_0 are model parameters.

The model has been constructed as the Monte-Carlo event generator PYQUEN (PYthia QUENched) and is available via Internet [5]. The event-by-event simulation procedure includes the generation of the initial parton spectra with PYTHIA [6] and production vertexes at given impact parameter, rescattering-by-rescattering simulation of the parton path length in a dense zone, radiative and collisional energy loss per rescattering, final hadronization with the Lund string model for hard partons and in-medium emitted gluons.

The full heavy ion event is simulated as a superposition of soft hydro-type state and hard multi-jets. The simple approximation [3] of hadronic liquid at “freeze-out” stage has been used to treat soft part of the event. Then the hard part of the event includes PYQUEN multi-jets generated according to the binomial distribution. The mean number of jets produced in AA events at a given impact parameter is a product of the number of binary NN sub-collisions and the integral cross section of hard process in pp collisions with the minimal transverse momentum transfer $p_T^{\min}$. The extended in such a way model has been also constructed as the fast Monte-Carlo event generator [7]. Note that ideologically similar approximation has been developed in [8].

3. Jet quenching at RHIC

In order to demonstrate the efficiency of the model, the jet quenching pattern in Au+Au collisions at RHIC was considered. Model parameters has been selected from the fit of η - and p_T - spectra at RHIC for different event centralities (see [3] for details). In particular, the nuclear modification of the hardest domain of p_T -spectrum was used to extract initial QGP conditions: $T_0 = 500$ MeV and $\tau_0 = 0.4$ fm/c. Figure 1 shows that our model reproduce p_T - and centrality dependences of nuclear modification factor R_{AA} (determined as a ratio of particle yields in AA and pp collisions normalized on the number of binary NN sub-collisions) for π_0 's measured by PHENIX [9] quite well. Another important tool to verify jet quenching is two-particle azimuthal correlation function $C(\Delta\varphi)$ – the distribution over an azimuthal angle of high- p_T hadrons in the event with $2 \text{ GeV}/c < p_T < p_T^{\text{trig}}$ relative to that for the hardest “trigger” particle with

$p_T^{\text{trig}} > 4 \text{ GeV}/c$. Figure 2 presents $C(\Delta\varphi)$ in pp and in central Au+Au collisions (data from STAR [10]). Clear peaks in pp collisions at $\Delta\varphi = 0$ and $\Delta\varphi = \pi$ indicate a typical dijet event topology. However, for central Au+Au collisions the peak near π disappears. It can be interpreted as the observation of monojet events due to the absorption of one of the jets in a dense medium. Figure 2 demonstrates that measured suppression of azimuthal back-to-back correlations is well reproduced by our model.




Figure 1. The nuclear modification factor R_{AA} for neutral pions in Au+Au collisions for two centrality sets. The points are PHENIX data, histograms are the model calculations.




Figure 2. The azimuthal two-particle correlation function for pp and for central Au+Au collisions. The points are STAR data, dashed and solid histograms are the model calculations for pp and Au+Au events respectively.

We leave beyond the scope of this paper the analysis of such important RHIC observables as the azimuthal anisotropy and particle ratios, which are sensitive to the soft physics. In order to study them, a more careful treatment of low- p_T particle production than our simple approach is needed [11] (the detailed description of space-time structure of freeze-out region, resonance decays, etc.).

4. Jet quenching at LHC

The developed model was applied to analyze various novel features of jet quenching at the LHC (see [12, 13, 14] for details). Let us just to enumerate some main issues.

Jets tagged by leading hadrons. The relation between in-medium softening jet fragmentation function (measured with leading hadrons) and suppression of jet rates due to energy loss out of jet cone, was analyzed in the work [12]. The specific anti-correlation between two effects allows one to probe parton energy loss mechanism (small-angular radiation versus wide-angular radiation and collisional loss).

Jets induced by heavy quarks. The possibility to observe B-jets tagged by an energetic muon in heavy ion collisions was analyzed in the work [13]. The significant softening b-jet fragmentation function due to b-quark energy loss is predicted.

Z/γ^ + jet production.* The channel with dimuon tagged jet production in heavy ion collisions was analyzed in the work [14]. We have found that the medium-induced partonic energy loss can result in significant P_T -imbalance between $\mu^+\mu^-$ pair and a leading particle in a jet, which is quite visible even for moderate loss.

5. Conclusions

The method to simulate jet quenching in heavy ion collisions has been developed. The model is the fast Monte-Carlo tool implemented to modify a standard PYTHIA jet event. The full heavy ion event is obtained as a superposition of a soft hydro-type state and hard multi-jets. The model is capable of reproducing main features of the jet quenching pattern at RHIC (the p_T -dependence of the nuclear modification factor and the suppression of azimuthal back-to-back correlations). The model was also applied to probe jet quenching in various new channels at LHC energy: jets tagged by leading particles, b-jets, dilepton-jet correlations. The further development of the model focusing on a more detailed description of low- p_T particle production is in the progress.

Acknowledgments

Discussions with A. Morsch, L.I. Malinina, C. Roland, L.I. Sarycheva and B. Wyslouch are gratefully acknowledged. I.L. thanks the organizers of “Quark Matter 2006” for the warm welcome and hospitality. This work is supported by grants N 04-02-16333 and 06-02-27355 of Russian Foundation for Basic Research and Contract N 02.434.11.7074 of Russian Ministry of Science and Education.

References

- [1] Baier R, Schiff D and Zakharov B G 2000 *Annual Rev. Nucl. Part. Sci.* **50** 37
- [2] Wang X-N 2004 *Phys. Lett.* **B 579** 299
- [3] Lokhtin I P and Snigirev A M 2006 *Eur. Phys. J.* **C 45** 211
- [4] Baier R, Dokshitzer Yu L, Mueller A H and Schiff D 1999 *Phys. Rev.* **C 60** 064902
- [5] <http://cern.ch/lokhtin/pyquen>
- [6] Sjostrand T 2001 *Comp. Phys. Com.* **135** 238
- [7] <http://cern.ch/lokhtin/hydro/hydjet.html>
- [8] Hirano T and Nara T 2004 *Nucl. Phys.* **A 743** 305
- [9] Adler S S *et al* (PHENIX Collab.) 2003 *Phys. Rev. Lett.* **91** 072301
- [10] Adler C *et al* (STAR Collaboration) 2002 *Phys. Rev. Lett.* **89** 202301
- [11] Amelin N S *et al* 2006 *Phys. Rev.* **C 74** 064901
- [12] Lokhtin I P and Snigirev A M 2003 *Phys. Lett.* **B 567** 39
- [13] Lokhtin I P, Sarycheva L I, Snigirev A M and Teplov K Yu 2004 *Eur. Phys. J.* **C 37**
- [14] Lokhtin I P, Sherstnev A V and Snigirev A M 2003 *Phys. Lett.* **B 599** 260